

Electron impact multiple ionization of C⁺, N⁺ and O⁺ ions.

J Lecointre^{1;5}, K A Kouzakov^{2;3}, D S Belic⁴, P Defrance⁵, Yu V Popov³, and V P Shevelko⁶

1. Haute Ecole Namur-Liège-Luxembourg HENALLUX, Département Ingénieur Industriel Pierrard, Rue d'Arlon 112, B-6760 Virton, Belgium.

2. Department of Nuclear Physics and Quantum Theory of Collisions, Faculty of Physics, Lomonosov Moscow State University, Moscow 119991, Russia.

3. Skobel'syn Institute of Nuclear Physics, Lomonosov Moscow State University, Moscow 119991, Russia.

4. University of Belgrade, Faculty of Physics, PO Box 386, 11000 Belgrade, Serbia.


5. Université Catholique de Louvain, Institute of Condensed Matter and Nanosciences, Chemin du Cyclotron 2, B-1348 Louvain-la-Neuve, Belgium.

6. Lebedev Physical Institute, Russian Academy of Sciences, Moscow 119991, Russia.

In the present study the animated crossed electron-ion beams method [1] is applied for measurement of absolute cross sections for electron impact single and multiple ionization of C⁺, N⁺ and O⁺ ions at incident electron energy values up to 2.5 keV.


The maximum cross sections for the multiply-charged products C^{q+} (q = 2 – 4) is found to range from 2.3×10⁻²⁰ cm² (for C⁴⁺) up to 6.3×10⁻¹⁷ cm² (for C²⁺); for N^{q+} (q = 2 – 5) they range from 3.0×10⁻²² cm² (for N⁵⁺) up to 5.1×10⁻¹⁷ cm² (for N²⁺) and, lastly, for O^{q+} (q = 2 – 5) they range from 5.5×10⁻²² cm² (for O⁵⁺) up to 5.2×10⁻¹⁷ cm² (for O²⁺).

The corresponding threshold energies are determined to satisfactorily compare to spectroscopic values.




Cross sections for electron impact double ionization of (a) C³⁺, (b) N³⁺ and (c) O³⁺: present experimental results (•), measurements of Zambra et al (×), measurements of Westermann et al (○), semi-empirical model of Talukder et al (---), and semi-empirical formula (—).

The Bethe-Fano plot show that for reasonably high energy of the incident electron, the quantity nE_i linearly depends on lnE_i, according to the Bethe formulae. The Born approximation in the interaction between a fast electron and an ion turns out to be sufficient already at impact energies of only about 1 keV.




Bethe-Fano plots for X^{q+} ions produced by electron impact ionization of (a) C⁺, (b) N⁺ and (c) O⁺: Straight lines result from the fitting procedure.



Cross sections for electron impact single ionization of (a) C⁺, (b) N⁺ and (c) O⁺: present experimental results (•), measurements of Yamada et al (○) and calculations based on the CBE approximation using the ATOM code (---).

The cross section values for single ionization reasonably agree with the calculations using the Coulomb-Born approximation with exchange [2]. Those for multiple ionization are found to compare well with the semi-empirical model for q = 3 [3], but they appear to be notably overestimated by a semi-empirical Bethe-Born type formula when q > 3 [4].



Ionization cross sections for C⁺, N⁺, O⁺ and Ne⁺ plotted as a function of the number of ejected electrons at E = 1995.1 eV: measurements of Lecointre et al (×) and present experimental results (open symbols) together with CBE calculations using the ATOM code for n = 1 and semi-empirical formulas [3] and [4] for, respectively, n = 2 and n = 3, 4 (filled symbols).

The remarkable point is that, at a given incident electron energy, the sequence of single and multiple ionization cross sections is observed to decrease exponentially with respect to the number of ejected electrons:

$$\sigma_n = \sigma_1 \exp\left[-(n-1)/q_0\right]$$

where σ_1 is the single ionization cross section and the fitting parameter q_0 appears to be an effective charge. This peculiarity is also observable for neutral targets, and therefore seems to be a general feature common to any atomic or ionic target, but, up to now, it seems to have not receive any qualitative or quantitative theoretical treatment.

References

- [1] P. Defrance et al 1981 *J. Phys. B: At. Mol. Phys.* **14** 103
- [2] V. P. Shevelko and L. A. Vainshtein 1993 *Atomic Physics for Hot Plasmas* (Bristol: Institute of Physics Publishing)
- [3] V. P. Shevelko et al 2005 *J. Phys. B: At. Mol. Opt. Phys.* **38** 525
- [4] V. P. Shevelko and H. Tawara 1995 *J. Phys. B: At. Mol. Opt. Phys.* **28** L589

Contact

julien.lecointre.henallux@gmail.com
kouzakov@gmail.com
popov@srd.sinp.msu.ru